



# CAPS 422-Mechanisms of Hypoxemia

Dr. Sally Osborne  
Department of Cellular & Physiological Sciences  
University of British Columbia  
Room 3602, D.H. Copp Building  
office: 604.822.3421  
email: [sally.osborne@ubc.ca](mailto:sally.osborne@ubc.ca)  
course website: [www.sallyosborne.com](http://www.sallyosborne.com)


# Important Terminology

Anoxia: absence of O<sub>2</sub> supply in the presence of perfusion- no oxygen

Asphyxia: absence of O<sub>2</sub> & accumulation of CO<sub>2</sub>.

Hypoxia: ↓ O<sub>2</sub> in the body, often specified where in the body  
e.g. tissue hypoxia, alveolar hypoxia

Hypoxemia: ↓ O<sub>2</sub> in the blood. Specifically, hypoxemia is determined by measuring the PO<sub>2</sub> of arterial blood (plasma)



## [P(A-a)O<sub>2</sub>]

Normal range=10-15 mmHg breathing room air, F<sub>I</sub>O<sub>2</sub>=0.21

- ▶ the normal range ↑ with age (1 mmHg per decade due to a ↓ PaO<sub>2</sub> as a result of ↑ ventilation perfusion mismatch)
- ▶ is due to venous admixture (anatomic shunt & ventilation perfusion mismatch in **health**)
- ▶ is due to venous admixture (anatomic shunt, ↑ ventilation perfusion mismatch & physiologic shunt in **disease states**)

# Causes of Hypoxemia


1. Hypoventilation
2. Low inspired oxygen
3. R-L shunt
4.  $\dot{V}/\dot{Q}$  inequality (a.k.a.  $\dot{V}/\dot{Q}$  mismatch)
5. Diffusion Impairment

Causes are subdivided into those with an increase in the  $P(A-a)O_2$  and those where the A-a gradient remains within the normal range.

# Hypoventilation

- $\uparrow$  PaCO<sub>2</sub> (hypercapnia)
- P(A-a)O<sub>2</sub> within normal range
- $\uparrow$  F<sub>I</sub>O<sub>2</sub> alleviates the hypoxemia
- mechanical ventilation required to eliminate hypercapnia

# Hypoventilation



## Causes:

1. Depression of CNS by drugs
2. Inflammation, trauma or hemorrhage in the brainstem
3. Abnormal spinal cord pathway
4. Disease of the motoneurons of the brain stem/spinal cord
5. Disease of the nerves supplying the respiratory muscles.
6. Disease of the neuromuscular junction
7. Disease of the respiratory muscles
8. Abnormality of the chest wall
9. Upper airway obstruction

## Low inspired oxygen ( $\downarrow$ PIO<sub>2</sub>)

- $\downarrow$  PIO<sub>2</sub> = (P<sub>B</sub> - 47 mmHg) FIO<sub>2</sub>
- P(A-a)O<sub>2</sub> within normal range
- $\downarrow$  PaCO<sub>2</sub> (hypocapnia due to hyperventilation in response to low arterial PO<sub>2</sub>)



# Right to Left Shunt

- $\uparrow P(A-a)O_2$
- $PaCO_2$  within the normal range

# Anatomic Shunt

**A portion of Blood bypasses the Lungs through an Anatomic Channel**

In all Healthy Individuals

- a portion of the bronchial circulation's venous blood drains into the pulmonary vein.
- a portion of the coronary circulation's venous blood drains through the thebesian veins into the left ventricle.


# Anatomic Shunt

A portion of Blood bypasses the Lungs through an Anatomic Channel

Disease States  
(Congenital abnormalities)

- intra-cardiac shunts
- intrapulmonary fistulas

- ❶ Ventricular Septal Defect
- ❷ Pulmonary Stenosis
- ❸ Hypertrophy of Rt. Ventricle
- ❹ Overriding Aorta




**Tetralogy of Fallot**

# Physiologic Shunt

A portion of cardiac output that goes through the normal pulmonary vasculature does not come into contact with alveolar air due to filling of the alveolar spaces with fluid


e.g.


- drowning
- pulmonary edema



## Key Clinical Feature of R-L Shunts

“the accompanying hypoxemia can not be corrected with supplemental oxygen”







# Ventilation Perfusion Inequality

- $\uparrow P(A-a)O_2$
- PaCO<sub>2</sub> within the normal range
- most common cause of hypoxemia in disease states




# Normal V/Q Inequality from the Apex to Base of the Lungs






"Shunt Like"  
 $\dot{V}/\dot{Q}=0$




Ideal Lung  
 $\dot{V}/\dot{Q}=1$



Dead Space  
 $\dot{V}/\dot{Q}=\text{infinity}$

zero ← low  $\dot{V}/\dot{Q}$  ← normal  $\dot{V}/\dot{Q}$  → high  $\dot{V}/\dot{Q}$  → infinity



$\downarrow O_2$     $\uparrow CO_2$   
 $\downarrow$   
 $\uparrow \dot{V}_E$


$\downarrow O_2$   
 $\uparrow CO_2$

$N O_2$   
 $N CO_2$

$\downarrow O_2, \uparrow CO_2$

$N O_2, N CO_2$

$\downarrow O_2, \uparrow CO_2$



# Diffusion Impairment

- $P(A-a)O_2$  normal at rest,  $\uparrow$ s with exercise
- $PaCO_2$  within the normal range
- a rare observation in clinical setting

Summary	arterial blood		venous blood		P(A-a)O <sub>2</sub>	Does supplemental oxygen (↑F <sub>I</sub> O <sub>2</sub> ) increase PaO <sub>2</sub> substantially?
	PO <sub>2</sub>	PCO <sub>2</sub>	PO <sub>2</sub>	PCO <sub>2</sub>		
<b>Hypoxemia</b>						
Hypoventilation	↓	↑	↓	↑	normal	yes
↓ P <sub>I</sub> O <sub>2</sub>	↓	↓	↓	↓	normal	yes
R-L Shunt	↓	normal	↓	normal	↑	no (depends on magnitude of the shunt)
Diffusion defect	↓	normal	↓	normal	↑ during exercise	yes
VA/Q inequality	↓	normal	↓	normal	↑	yes
<b>Tissue hypoxia</b>						
Anemic hypoxia	normal	normal	↓	normal	normal	no
CO poisoning	normal	normal	↓	normal	normal	possibly
Stagnant hypoxia	normal	normal	↓	normal	normal	no
Histotoxic hypoxia	normal	normal	↑	normal	normal	no

**Important note** that mixed causes of hypoxemia occur frequently. It is often impossible to define the extent of the contribution of each mechanism in the acutely ill patient.